

## Deliverable 1.1

### Inventory Report

Authors and affiliation:

**Melanie Witthöft**  
**Hans Doornenbal**

[LBEG, TNO]

E-mail of lead author:

**Melanie.witthoeft@lbeg.niedersachsen.de**

Version: 19-12-2018

This report is part of a project that has received funding by the European Union's Horizon 2020 research and innovation programme under grant agreement number 731166.



Deliverable Data		
<b>Deliverable number</b>	D1.1	
<b>Dissemination level</b>	Public	
<b>Deliverable name</b>	Inventory report	
<b>Work package</b>	WP1, Cenozoic geothermal reservoirs in the northern onshore Dutch-German cross border region	
<b>Lead WP/Deliverable beneficiary</b>	LBEG (Lower Saxony, Germany)	
Deliverable status		
<b>Submitted (Author(s))</b>	18/12/2018	Melanie Witthöft, Hans Doornenbal
<b>Verified (WP leader)</b>	18/12/2018	Melanie Witthöft
<b>Approved (Project leader)</b>	19/12/2018	Stefan Knopf

## GENERAL INTRODUCTION

In WP1 of 3DGEU Lower Saxony and the Netherlands are involved. Lower Saxony is represented by the Authority for Mining, Energy and Geology (Landesamt für Bergbau, Energie und Geologie, LBEG) and the Netherlands are represented by the Netherlands Organisation for Applied Scientific Research (TNO). Both institutions are experienced in three dimensional modeling of the subsurface.

In 3DGEU both partners will work in a cross border pilot area (see Figure 1) on harmonization of existing 3D models and Cenozoic geothermal reservoir maps.



Figure 1: Overview of the pilot area: red box in the left picture marks the participating countries, the blue area in the right picture shows the cross border pilot area in the Netherlands and Lower Saxony (Eurostat, 2018).

This report (deliverable of task 1.1) is a state of the art inventory of existing 3d models (see chapter 1) and Cenozoic reservoir maps and their properties (see chapter 2) in Lower Saxony and the Netherlands.

This inventory will be the base for the following cross border tasks in this project:

- In task 1.2 LBEG and TNO strive to establish a consistent 3d model of the pilot area. Therefore we present in chapter 3 criteria for harmonization.
- Harmonized distribution, depth and thickness maps of Cenozoic geothermal reservoirs will be the result of task 1.3.
- Properties like porosity, permeability and thermal conductivity will be gathered and loaded into a database in task 1.4, which will be the base to create Cenozoic geothermal maps under task 1.5.
- In task 1.6 a harmonized map of the hydraulic barrier between fresh groundwater and the deep salt groundwater system is presented as a decision support tool for planners.



## TABLE OF CONTENTS

1	EXISTING 3D MODELS.....	2
1.1	Involved countries .....	2
1.2	Extent.....	3
1.3	Input data .....	3
1.4	Date of publication .....	4
1.5	Model type.....	5
1.6	Stratigraphical Nomenclature.....	5
1.7	Modeled layers .....	8
1.8	Faults.....	13
1.9	Publications.....	14
1.10	Modeling software.....	14
1.11	Confidentiality and availability .....	14
1.12	Output data (sort/type, format) .....	15
1.13	Coordinate system.....	15
1.14	Height reference system.....	15
2	EXISTING MAPS OF CENOZOIC RESERVOIR ROCKS .....	16
2.1	Input data .....	17
2.2	Output data (sort/type, format) .....	17
2.3	Confidentiality and availability .....	18
3	HARMONIZATION.....	19
3.1	Harmonized main Mesozoic stratigraphic units .....	19
3.2	Harmonising Cenozoic stratigraphic units .....	19
3.3	Faults.....	23
4	LIST OF REFERENCES.....	24



## 1 EXISTING 3D MODELS

Our work is based on four 3D models: Geotektonischer Atlas 3D (GTA3D), Digitaal Geologisch Model diep (DGM-deep), Digitaal Geologisch Model Noord-Nederland (DGM-NNL) and Southern Permian Basin Atlas (SPBA). In this report the acronyms as mentioned in table 1, will be used.

Table 1: Overview of existing 3D models and used acronyms in this report.

3D models	
acronym	name
GTA3D (LBEG)	Geotektonischer Atlas 3D
DGM-deep V5.0 (TNO)	Digitaal Geologisch Model: DGM- diep
DGM-NNL (TNO)	Digitaal Geologisch Model Noord-Nederland
SPBA	Southern Permian Basin Atlas

Elaboration on criteria for harmonization of 3D models requires the knowledge of differences between the datasets. The properties and characteristics of these models will be listed in this chapter mainly as tables to derive criteria which will be used for the harmonization (chapter 3).

### 1.1 Involved countries

The involved countries, who have worked on the four 3D models are listed in table 2.

Table 2: Overview of involved nations.

3D model	Involved countries
GTA3D	Lower Saxony (Germany)
DGM-deep V5.0	The Netherlands
DGM-NNL	The Netherlands
SPBA	United Kingdom, Belgium, The Netherlands, Denmark, Germany and Poland

## 1.2 Extent

The extent of each 3D model is shown in figure 2. GTA3D and DGM-deep V5.0 extend over the whole country area of Lower Saxony respectively the Netherlands. DGM-NNL extends over the northeastern part of the Netherlands (see Figure 2). Only SPBA is a crossborder model with more countries involved.




Figure 2: 1) Extent of GTA3D model (grey shadow) (LBEG, 2018), 2) Extent of DGM-deep V5.0 (blue border), 3) Extent of DGM NNL (blue border) and 4) Extent of SPBA model (blue border) (Doornenbal et al., 2010).

## 1.3 Input data

Table 3: Different input data of existing 3D models.

3D model	Input data
GTA3D	14 depthmaps of the Tectonic Atlas of Northwest Germany and the German North Sea Sector (Baldschuhn et al., 2001); depth of Quaternary basis 1: 500 000 (NLfB, 1995); digital surface model; state of knowledge: ~ 2001
DGM-deep V5.0	seismic data and well data
DGM-NNL	3D seismic data and well data
SPBA	seismic data and well data

Database of the models is quite different at first glance. GTA3D is a model established on an older seismic and geological interpretation, executed before 1996. It is a model based on structural depth maps of the 'Geotektonischer Atlas von Nordwest-Deutschland und dem deutschen Nordsee-Sektor' (Baldschuhn et al., 2001). These maps have been generated by interpretation of 2D seismic data and

well information. 2D geological cross sections, thickness maps and paleotectonic maps complete these maps. Based on digitized structural maps, triangulated three dimensional surfaces were created to represent the base of the horizons (see Figure 3). For the other 3D models seismic data and well data was used as direct input data.




Figure 3: GTA3D – example of the modeling workflow of the Lower Cretaceous horizon, based on digitized maps (Bombien et al. 2012).

In GTA3D, data inconsistencies, caused by intersection of different structural maps in 3D model, have not been revised. Figure 4 shows an example of intersecting base surfaces. Due to 2-dimensional mapping of the input data, which was done in the 1980's and 1990's without the use of 3D modeling software, these inconsistencies mainly occur in areas of complex fault systems.




Figure 4: GTA3D - example of data inconsistencies; intersection of base surface of Middle Jurassic (blue) and base surface of Upper Jurassic+Wealden (green) (Helms 2018, unpublished).

## 1.4 Date of publication

It is necessary to record the date of publication. Table 4 lists the date of publication of the 3D models.

Table 4: Date of first publication.

3D model	Date of publication
GTA3D	17.07.2012
DGM-deep V5.0	Q1 2019
DGM-NNL	Q1 2019
SPBA	June 2010



## 1.5 Model type

Table 5: Different modeltypes of existing 3D models.

3D model	Model type
GTA3D	regional, main lithostratigraphical intervals
DGM-deep V5.0	regional, main lithostratigraphical intervals
DGM-NNL	detailed, reservoir layers
SPBA	regional, main lithostratigraphical intervals

## 1.6 Stratigraphical Nomenclature

In general 3D models exist of lithostratigraphic layers which are described in Stratigraphical Nomenclatures (see table 6).

Table 6: Different Stratigraphical Nomenclatures of existing 3D models.

3D model	Stratigraphical Nomenclature
GTA3D	lithostratigraphic units (see Table 7)
DGM-deep V5.0	<a href="https://www.dinoloket.nl/nomenclator">https://www.dinoloket.nl/nomenclator</a> and see Figure 5
DGM-NNL	<a href="https://www.dinoloket.nl/nomenclator">https://www.dinoloket.nl/nomenclator</a> and see Figure 5
SPBA	Each of the participating geological surveys has published a lithostratigraphic nomenclature for their country: UK (Cameron et al., 1993; Johnson et al., 1994; Lott & Knox, 1994; Waters et al., 2007), Belgium (Bultynck & Dejonghe, 2002), the Netherlands (Van Adrichem Boogaert & Kouwe, 1993), Denmark (Michelson et al., 2003; Schiøler et al., 2005), Germany (STD 2002; Menning & Hendrich, 2005), and Poland (Wagner, 2008).

Table 7: Stratigraphic terms in northwestern part of Germany, Age in million years (Ma) modified after (Baldschuhn et al., 2001) based on (Menning, 1997).

Age of base	Epoch (german)	Stratigraphic Unit (german)	Code (german)	Stratigraphic Series or stage (english)
1,8 Ma	Quartär	Quartär	(q)	Pleistocene-Holocene
24 Ma	Neogen (Jungtertiär)	Pliozän	(tpl)	Pliocene
		Miozän	(tmi)	Miocene
65 Ma	Paläogen Alttertiär	Oligozän	(tol)	Oligocene
		Eozän	(teo)	Eocene
		Oberpaläozän	(tpao)	Upper Paleocene
		Dan	(td)	Danian
99 Ma	Oberkreide	Maastricht	(krma)	Maastrichtian
		Campán	(krca)	Campanian
		Santon	(ksra)	Santonian
		Coniac	(krcc)	Coniacian
		Turon	(krt)	Turonian
		Cenoman	(krc)	Cenomaian
continued on next page				



Age of base	Epoch (german)	Stratigraphic Unit (german)	Code (german)	Stratigraphic Series or stage (english)	
144 Ma	Unterkreide	Alb	(krl)	Albian	
		Apt	(krp)	Aptian	
		Barrême	(krb)	Barremian	
		Hauterive	(krh)	Hauterivian	
		Valangin	(krv)	Valanginian	
		Berrias = Wealden	(Wd)	Berriasiyan	
159 Ma	Oberjura (Malm)	Serpulit	(joS)	Tithonian	
		Münster Mergel	(joM)	Tithonian	
		Eimbeckhäuser P.-K.	(joE)	Tithonian	
		Gigas-Schichten	(joG)	Tithonian	
		Kimmeridge	(joKI)	Kimmeridgian	
		Korallenoolith	(joK)	Oxfordian	
		Heersumer Sch.	(joH)	Oxfordian	
180 Ma	Dogger (Mittlerer Jura)	Callovium	(jmcl)	Callovian	
		Bathonium	(jmbl)	Bathonian	
		Bajocium	(jmbj)	Bajocian	
		Aalenium	(jmal)	Aalenian	
206 Ma	Lias (Unterer Jura)	Toarcium	(jutc)	Toarcian	
		Pliensbachium	(jupl)	Pliensbachian	
		Sinemurium	(jusi)	Sinemurian	
		Hettangium	(juhe)	Hettangian	
231 Ma	Keuper	Rhät	(ko)	Rhaetian	
		Steinmergelkeuper	(km4)	Norian	
		Oberer Gipskeuper	(km3)	Carnian	
		Schliffsandstein	(km2)	Carnian	
		Unterer Gipskeuper	(km1)	Ladinian	
		Lettenkeuper	(ku)	Ladinian	
240 Ma	Muschelkalk	Ob. Muschelkalk	(mo)	Ladinian-Ansian	
		Mittl. Muschelkalk	(mm)	Ansian	
		Unt. Muschelkalk	(mu)	Ansian	
251 Ma	Buntsandstein	Ob.Buntsandstein	Röt	(so)	Ansian
			Solling-Folge	(smS)	Olenekian
		Mittl. Buntsandstein	Hardegsen-Folge	(smH)	Olenekian
			Detfuth-Folge	(smD)	Olenekian
			Volpriehausen-Folge	(smV)	Olenekian
		Unt. Buntsandstein	Quickborn-Folge	(smQ)	Olenekian
			Bernburg-Folge	(suB)	Induan
			Calvörde-Folge	(suC)	Induan
258 Ma	Zechstein		Mölln-Zyklus	(z7)	Changhsingian
			Friesland-Zyklus	(z6)	Changhsingian
			Ohre-Zyklus	(z5)	Changhsingian
			Aller-Zyklus	(z4)	Changhsingian
			Leine-Zyklus	(z3)	Wuchiapingian
			Staßfurt-Zyklus	(z2)	Wuchiapingian
			Werra-Zyklus	(z1)	Wuchiapingian
300 Ma	Rotliegend		Oberrotliegend	(ro)	Artiskian-Wuchiapingian
			Unterrotliegend	(ru)	Gzhelian-Artiskian
326,3 Ma	Oberkarbon		Stefan	(cst)	Kasimovian-Gzhelian
			Westfal	(cw)	Bashkirian-Kasimovian
			Namur	(cn)	Serpukhovian-Bashkirian
353,8 Ma	Unterkarbon	Dinant	(cd)	Tournaisian-Serpukhovian	




Figure 5: Stratigraphical Nomenclature used in DGM-deep V5.0 and DGM-NNL (TNO, 2018).



## 1.7 Modeled layers

Table 8: Different modeled layers of existing 3D models.

3D model	Modeled layers
GTA3D	15 layers (see Table 9 and Figure 6)
DGM-deep V5.0	NU, N, CK, KN, S, ATPO, AT, RN, RB, ZE, RO + 2 Carb.horizons (see Table 10)
DGM-NNL	67 layers (see Table 10)
SPBA	regional, main lithostratigraphical intervals (see Figure 7 and Figure 8)

Table 9: 14 lithostratigraphic units published in GTA; in GTA3D Quaternary base was additionally modeled.

Stratigraphic Units Geotectonic Atlas NW-Germany	Acronym
Base deepest Middle Miocene (Reinbeck)(NN5) to highest Pliocene	03_tmim-tpl
Base deepest Lower Miocene (Vierlande) (NN1) to youngest Lower Miocene (Hemmoor) (NN4)	04_tmiu
Base Middle Oligocene (Rupel) (NP23) to youngest Upper Oligocene (Neochatt) (NP25)	05_tolm-tolo
Base Brussels Sand to Top Lower Oligocene	06_tolm-teoo
Base Upper Paleocene (Landen) (NP8) to youngest Lower Eocene (Lower Eocene 4) (NP13)	07_tpao-teou
Base Cenoman to highest Dan	08_kro
Base deepest marine Lower Cretaceous (Valangin) to highest Lower Cretaceous (Oberalb)	09_kru
Base deepest Upper Jurassic (Oxford) to "Wealden" (Berrias)	10_jo-Wd
Base highest Lias (Obertoarcium) to highest Dogger (Callovium)	11_jutco-jmclo
Base lowest Lias (Hettangium) to Upper Lias (Lower Toarcium, Posidonia Shale)	12_juhe-jutcu
Base deepest Keuper (Lettenkeuper) to Upper Keuper (Upper Rhaetian)	13_k
Base Upper Buntsandstein (Roetsalinar) to Upper Muschelkalk (Ceratitenschichten)	14_so-m
Base deepest Lower Buntsandstein (Calvoerde) to highest Middle Buntsandstein (Solling)	15_su-sm
Base deepest Zechstein (Kupferschiefer Werra Folge) (T1) to highest Zechstein (Moelln-Cycle) (z7) oder Broeckelschiefer	16_z

Geologische Einheiten	System/Periode	AraName/Ara	Alter (Ma) (DSK 2002)
Quärtär ungegliedert	Quärtär		1,8 – 0
Mittelmiozän bis Pliozän			16,5 – 1,8
Untermiozän			23,8 – 16,5
Rupel bis Oberoligozän	Tertiär	Kanozoikum	32,5 – 23,8
Mitteldorfazän bis Obereozän			49 – 32,5
Oberpaläozän bis Untereozän			58 – 49
Oberkreide	Kreide		99 – 65
Marine Unterkreide			142 – 99
Oberjura und "Wealden"	Jura/Kreide		156,5 – 137
Mitteljura (Dogger)	Jura	Mesozoikum	178 – 156,5
Unterjura (Lias)			200 – 178
Keuper			235 – 200
Röt und Muschelkalk	Trias		244,5 – 235
Unterer und Mittlerer Buntsandstein			251 – 244,5
Zechstein	Perm	Paläozoikum	258 – 251

Figure 6: General legend of geological units, age and colours used in 3dD model GTAD3D (LBEG, 2008).




Table 10: Stratigraphic units and model layers of DGM-deep V5.0 and DGM-NNL (Ten Veen et.al., in prep).

Quaternary	Naaldwijk	Early and Middle Jurassic	Werkendam Formation
	Boxtel		Aalburg Formation
	Eem		<b>SLEEN FORMATION (BASE AT)</b>
	Drente		Upper Keuper Claystone
	Drachten		Dolomitic Keuper
	Urk Tynje		Red Keuper Claystone
	Peelo		Red Keuper Evaporite
	Urk		Middle Keuper Claystone
	Appelscha		Main Keuper Evaporite
Neogene	Peize		Lower Keuper Claystone
	Maassluis		Upper Muschelkalk
	Oosterhout		Middle Muschelkalk Marl
	<b>BREDA (BASE NU)</b>		Muschelkalk Evaporite
Paleogene	Rupel Clay	Triassic	Lower Muschelkalk
	<b>VESSEM MEMBER</b>		Upper Röt Claystone
	<b>ASSE MEMBER</b>		Upper Röt Evaporite
	<b>BRUSSEL MEMBER</b>		Intermediate Röt Claystone
	<b>IEPER MEMBER</b>		Main Röt Evaporite
Late Cretaceous	Basal Dongen Tuffite		<b>SOLLING CLAYSTONE (BASE RN)</b>
	<b>LANDEN CLAY (BASE N)</b>		Basal Solling Sandstone
	Ommelanden Formation		Hardegsen Formation
Early Cretaceous	Plenus Marl		Defurth Claystone
	<b>TEXEL FORMATION (BASE CK)</b>		Lower Defurth Sandstone
	Upper Holland Marl		Volpriehausen Clay-Siltstone
	Middle Holland Claystone		Lower Volpriehausen Sandstone
	Lower Holland Marl		Rogenstein
Late Jurassic	Vlieland Claystone I	Permian	<b>MAIN CLAYSTONE (BASE RB)</b>
	Friesland Member		<b>ZECHSTEIN GROUP (BASE ZE)</b>
	Vlieland Claystone II		
	<b>BENTHEIM SANDSTONE (BASE KN)</b>		
	Upper Coevorden		
	Middle Coevorden		
	Lower Coevorden		
	Serpulite Member		
	Weiteveen Upper Marl		
	Weiteveen Upper Evaporite		
	Weiteveen Lower Marl		
	Weiteveen Lower Evaporite		
<b>WEITEVEEN BASAL CLASTICS (BASE S)</b>			



Io. Name of mapped lithostratigraphic horizon	Description	Approximate age (Ma)
1 Near base Tertiary	Top Danian or top of the Chalk Group	61
2 Base Upper Cretaceous	Base of the Chalk Group (base Cenomanian)	100
3 Near base Lower Cretaceous	Approximately near base Ryazanian; may correspond to an unconformity	140
4 Near base Upper Jurassic	This horizon corresponds to a level varying from the Callovian to the base of the Oxfordian.	162
5 Near base Middle Jurassic	This level corresponds to the top of the Lias Group (base West Sole Group and equivalents in the UK; top of the Posidonia Shale Formation in the Netherlands) and is a marked unconformity over much of the area	176
6 Near base Lower Jurassic	Base of the Lias Group. This level corresponds to the mudstone/sandstone boundary in the upper part of the Rhaetian, which is often an unconformity	201
7 Near base Upper Triassic	This level corresponds to a change from clastic-dominated to limestone-dominated sediments. It is not mapped in the UK and Denmark	237
8 Near base Middle Triassic	Base of the Röt evaporites	245
9 Near base Lower Triassic	Base of the Buntsandstein	253
0 Base Zechstein	The base-Kupferschiefer reflector is mostly indistinct or not present because of low impedance contrast. In general, the seismic reflector that has been used is a younger horizon: top basal anhydrite (Stassfurt series). An agreed thickness has been added to this marker to create the base-Zechstein map	258
1 Base upper Rotliegend	This level corresponds with the base of the upper Rotliegend clastics. In general, the horizon was created by combining the upper Rotliegend thickness map, based on well data, and the depth of the base-Zechstein horizon	265

Figure 7: Overview of the lithostratigraphic horizons mapped in the SPBA area. These horizons have been numbered from 1 to 11 and are indicated in Figure 8 (Doornenbal et al., 2010).






Figure 8: Stratigraphical correlation chart as used for SPBA; red lines and numbers show the (seismically mapped and) modeled layers (Doornenbal et al., 2010).

## 1.8 Faults

Information and data about faults and fault systems have been treated differently in the 3D models (see Table 11).

Table 11: Treatment of fault information in 3D models.

3D model	Representation of faults
GTA3D	no fault surfaces or correlation of fault traces were included; the vertical displacement on the stratigraphic horizons show indirectly the existence of faults.
DGM-deep V5.0	Fault surfaces have been mapped in seismics and partly modeled in 3D
DGM-NNL	Fault surfaces have been mapped in seismics and modeled in detail in 3D
SPBA	Main fault traces were retrieved from existing depth maps

In GTA3D, faults are not shown as fault surfaces. Information about faults in original GTA maps (see Figure 9) were transferred into the model as vertical displacements within the stratigraphic horizons.




Figure 9: Detail of GTA map "base Lower Jurassic" showing fault traces and additional information (e.g., dip angle and dip direction and the fault shows a horizontal and vertical displacement) (e.g., red circle) of faults; modified after (Baldschuhn et al. 2001).

In the models DGM-deep and DGM-NNL faults have been mapped during the seismic interpretation. In the next step, the faults have been modelled in 3D by using the Petrel software. In the SPBA model the main faults have been retrieved from existing depth maps.



## 1.9 Publications

Table 12: Publication list.

3D model	Publications
GTA3D	Bombien et al. (2012) Der Geotektonische Atlas von Niedersachsen und dem deutschen Nordseesektor als geologisches 3D-Modell. In: Geowissenschaftliche Mitteilungen : GMit / Berufsverband Deutscher Geowissenschaftler.- 48 (2012), Seite 6-13.
DGM-deep V5.0	Kombrink, H., Doornenbal, J.C., Duin, E.J.T., den Dulk, M., van Gessel, S.F., ten Veen, J.H. & Witmans, N. (2012) New insights into the geological structure of the Netherlands; results of a detailed mapping project. Netherlands Journal of Geoscience 91-4: 419-446.
DGM-NNL	Ten Veen, J.H., Kruisselbrink, A.F., den Dulk, M. and Witmans, N., in prep. Digital Geological Model Noord Nederland (DGM-NNL), TNO report, xx pp.
SPBA	Doornenbal, J.C. and Stevenson, A.G. (editors), 2010. Petroleum Geological Atlas of the Southern Permian Basin Area. EAGE Publications b.v. (Houten). ISBN 978-90-73781-61-0

## 1.10 Modeling software

Table 13: Used 3D modeling software for generating existing 3D models.

3D model	Modeling software
GTA3D	Paradigm Gocad
DGM-deep V5.0	Petrel
DGM-NNL	Petrel
SPBA	Petrel

Different modeling software was used. While Lower Saxony modeled with Paradigm Gocad, the Netherlands used Petrel software. GTA3D consists of triangulated surfaces (ts-files) and DGM-deep, DGM-NNL and SPBA were modeled as grids.

## 1.11 Confidentiality and availability

Table 14: Confidentiality of existing 3d models and information about data availability.

3D model	Confidentiality and availability
GTA3D	public on <a href="http://nibis.lbeg.de/cardomap3/?TH=1410">http://nibis.lbeg.de/cardomap3/?TH=1410</a> as 3dpdf or browser version
DGM-deep V5.0	public on <a href="http://www.nlog.nl">www.nlog.nl</a> and <a href="http://www.dinoloket.nl">www.dinoloket.nl</a> , Q1 2019
DGM-NNL	public, Q1 2019
SPBA	Atlas public via <a href="http://www.nlog.nl">www.nlog.nl</a> , GIS product (maps+databases) for sale



## 1.12 Output data (sort/type, format)

Table 15: Description of output data.

3D model	Output data
GTA3D	3D pdf (scale 1: 100 000) or GST-browser version; Gocad-t-surf-Format; cross sections and artificial well information on <a href="http://nibis.lbeg.de/cardomap3/">http://nibis.lbeg.de/cardomap3/</a>
DGM-deep V5.0	twt-, isopach-, depth-, V0-maps in pdf, arc-grid and zmap-grid, see <a href="https://www.nlog.nl/en/dgm-deep-v4-onshore">https://www.nlog.nl/en/dgm-deep-v4-onshore</a>
DGM-NNL	twt-, isopach-, depth maps
SPBA	Atlas on paper and pdf; GIS-product on DVD

## 1.13 Coordinate system

Table 16: Overview of coordinate system of existing 3D models.

3D model	Coordinate system
GTA3D	ETRS89 / UTM zone 32N (zE-N) (Europäisches Terrestrisches Referenzsystem von 1989 / Universale Transversale Mercatorabbildung Zone 32N) (EPSG 4647)
DGM-deep V5.0	Amersfoort/RD New, spheroid: Bessel 1841 (EPSG 28992)
DGM-NNL	Amersfoort/RD New, spheroid: Bessel 1841 (EPSG 28992)
SPBA	Lambert conformal conical (EPSG 5643)

## 1.14 Height reference system

Table 17: Overview of height reference system of existing 3D models

3D model	Height reference system
GTA3D	DE_DHHN92_NH
DGM-deep V5.0	onshore: Actueel Hoogtebestand Nederland v2 (AHN2) with reference level NAP (Nieuw Amsterdams Peil; offshore: msl (mid sea level)
DGM-NNL	Actueel Hoogtebestand Nederland v2 (AHN2) with reference level NAP (Nieuw Amsterdams Peil)
SPBA	msl ( mid sea level)



## 2 EXISTING MAPS OF CENOZOIC RESERVOIR ROCKS

In this chapter an inventory of existing maps of Cenozoic reservoir rocks of the study area is described.

In the Netherlands a public, web-based geographical information system "ThermoGIS V2.0" has been developed with the main goal of supporting companies and the government to develop geothermal energy in the Netherlands (see <https://www.thermogis.nl/en>). ThermoGIS provides depth, thickness, porosity and permeability maps of many potential aquifers in the Netherlands. Geothermal performance maps are calculated with the use of an integrated, stochastic, techno-economic performance module. The most important outputs of ThermoGIS are geothermal potential maps of the Netherlands that can be viewed in a map viewer. For our study area ThermoGIS provides two important aquifers i.e. Vessem Member (NMRFV) and the Brussels Sand Member (NLFFS), which are also geothermal reservoirs.

In Lower Saxony average thermal conductivity maps provide non-interpolated point information, which is based on well data. Average values per well are available for depths intervals from ground level to 40 m, 60 m, 80 m and 100 m depth. For our study area these maps show information mostly for quaternary deposits, which mostly consist of quaternary aquifers and in deeper parts of tertiary aquitards.

These existing maps will not be used in this project. The deliverable of task 1.5 will be geothermal maps for the pilot area based on our modeling work.

Main modeled Cenozoic layers such as top Neogene, base Neogene and base Paleogene from the existing models GTA3D, DGM-deep and DGM-NNL will be harmonized. Subsequently by using these layers as a geometrical reference together with the geometry (top, depth, thickness) from public wells the geometrical (top, base, thickness) maps of the geothermal reservoirs will be generated for the Cenozoic subsurface in the pilot area.



## 2.1 Input data

Table 18: different input data of existing maps

map	input data
Thermal conductivity for vertical heat exchangers (reference depth to 40m)	well information from ground level up to 40m depth, assessment of thermal conductivity (after VDI (2010), measurements and Ad-Hoc AG Hydrogeologie (2008), groundwatersurface
Thermal conductivity for vertical heat exchangers (reference depth to 60m)	well information from ground level up to 60m depth, assessment of thermal conductivity (after VDI (2010), measurements and Ad-Hoc AG Hydrogeologie (2008), groundwatersurface
Thermal conductivity for vertical heat exchangers (reference depth to 80m)	well information from ground level up to 80m depth, assessment of thermal conductivity (after VDI (2010), measurements and Ad-Hoc AG Hydrogeologie (2008), groundwatersurface
Thermal conductivity for vertical heat exchangers (reference depth to 100m)	well information from ground level up to 100m depth, assessment of thermal conductivity (after VDI (2010), measurements and Ad-Hoc AG Hydrogeologie (2008), groundwatersurface
ThermoGIS 2.0 - NLFFS – Brussels sand Member	Some harmonized main modeled Cenozoic layers together with the geometry (top, depth, thickness) from wells
ThermoGIS 2.0 – NMRFV – Vessem Member	Some harmonized main modeled Cenozoic layers together with the geometry (top, depth, thickness) from wells

## 2.2 Output data (sort/type, format)

Table 19: description of output data of existing maps

map	output data
Thermal conductivity for vertical heat exchangers (reference depth 40m)	point information with additional depth information
Thermal conductivity for vertical heat exchangers (reference depth 60m)	point information with additional depth information
Thermal conductivity for vertical heat exchangers (reference depth 80m)	point information with additional depth information
Thermal conductivity for vertical heat exchangers (reference depth 100m)	point information with additional depth information
ThermoGIS 2.0 - NLFFS – Brussels sand Member	Top, base, thickness, porosity and permeability maps as grid+pdf
ThermoGIS 2.0 – NMRFV – Vessem Member	Top, base, thickness, porosity and permeability



	maps as grid+pdf
--	------------------

## 2.3 Confidentiality and availability

Table 20: status of confidentiality and link to the output data

map	Confidentiality and availability
Thermal conductivity for vertical heat exchangers (reference depth to 40m)	public under <a href="http://nibis.lbeg.de/cardomap3/">http://nibis.lbeg.de/cardomap3/</a>
Thermal conductivity for vertical heat exchangers (reference depth to 60m)	public under <a href="http://nibis.lbeg.de/cardomap3/">http://nibis.lbeg.de/cardomap3/</a>
Thermal conductivity for vertical heat exchangers (reference depth to 80m)	public under <a href="http://nibis.lbeg.de/cardomap3/">http://nibis.lbeg.de/cardomap3/</a>
Thermal conductivity for vertical heat exchangers (reference depth to 100m)	public under <a href="http://nibis.lbeg.de/cardomap3/">http://nibis.lbeg.de/cardomap3/</a>
ThermoGIS 2.0 - NLFFS – Brussels sand Member	Public under <a href="https://www.thermogis.nl/en/map-viewer">https://www.thermogis.nl/en/map-viewer</a>
ThermoGIS 2.0 - NMRFV – Vessem Member	Public under <a href="https://www.thermogis.nl/en/map-viewer">https://www.thermogis.nl/en/map-viewer</a>



## 3

## HARMONIZATION

For compilation of existing 3D models well and seismic data have been used. The decision, which layers have been modeled was derived on base of the input data. Therefore seismostratigraphic or lithostratigraphic horizons may be the base for the different models. In January 2019 we start to strive the harmonization of existing 3D models to a consistent 3D model for several Mesozoic main intervals (chapter 3.1) and some Cenozoic intervals (chapter 3.2). During this work we will examine for which of these four 3D models a harmonization is possible and in which range. Detailed inventory of modeled layers will be the basis for the following work. Realisation of harmonisation is only possible, if it is geological reasonable. Otherwise modeled layers are completely different and not comparable or traceable in other models.

In chapter 1 the differences between the four 3D models have been described. For elaborating a harmonized 3D model it is important to agree first on a common coordinate system for our pilot area to prevent inaccuracy and mistakes in geographic position.

### 3.1 Harmonized main Mesozoic stratigraphic units

During the SPBA project the seismic interpreted main horizons in the different countries have been already compared, also for Germany and the Netherlands (GTA3D and DGM-deep, see Figure 8). For the Mesozoic a total of eight horizons have been mapped, from base Triassic to base Cenozoic (see Figure 11). For three horizons, i.e. near Base Upper Jurassic, near base Middle Jurassic and near Base Lower Jurassic, the seismic interpreted horizons were picked on different stratigraphic positions on both sides of the German-Dutch border (see Figure 8). For each of these three horizons we have to determine the difference in age and in depth, afterwards we have to decide how to solve the differences.

### 3.2 Harmonising Cenozoic stratigraphic units

GTA3D depicts the Cenozoic layers in more detail with six stratigraphic layers: see table 9. DGM-NNL depicts Cenozoic layers more in detail: see table 10. DGM-deep depicts two Cenozoic horizons: base Upper North Sea Group (near base Neogene) and base Lower North Sea Group (near base Paleogene). In Figure 10 the seismic mapped horizons for Cenozoic are indicated, which have been modelled in the GTA3D (Germany) and DGM-NNL (Northeast-Netherlands) models. From this figure we could conclude that the following four horizons were seismically mapped on approximately the same level in both models GTA3D and DGM-NNL:

- base Landen Clay Member (NLLFC = 07 tpa0-teou),
- base Brussels Sand Member (NLFFS=06 tolm-teoo),
- base Rupel Formation (NMRF= 05 tolm-tolo) and
- base Quaternary.

Only for the seismically mapped base Upper North Sea Group (=NUBA) in DGM-deep or DGM-NNL it should be decided whether it will be possible to harmonise this level with "04 tmiu" or "03 tmim-tpl" in the GTA3D model. The five mentioned horizons above could be harmonized along the German-Dutch border, only on the Dutch side of the pilot area some horizons should still be seismically mapped in the area outside the DGM-NNL area. Subsequently by combining these main horizons as a geometrical reference together with the geometry (top, depth, thickness) from public wells the



---

geometrical maps (top, base, thickness) of the geothermal reservoirs will be generated for the Cenozoic (except Holocene) subsurface in the pilot area.

The geometrical maps of these geothermal reservoirs (aquifers) will be the base for the geothermal property maps, which will be created in Task 1.5.




Figure 10: Tectonostratigraphic Chart for Cenozoic (modified after (Doornenbal and Stevenson, 2010)). The red line is the seismically mapped near base Cenozoic in the SPBA-project. Blue lines show the seismically mapped horizons of DGM-NNL (Rupel Formation consists of modeled layers: Rupel Clay and Vessem Member) and the green lines show the seismically mapped horizons of GTA3D.




Figure 11: correlation of modelled layers for GTA3D (red lines), DGM deep (blue lines) and SPBA ( green lines) with names of the modeled horizons(modified after (Kombrink et al., 2012))

### 3.3 Faults

As described in Chapter 1.8 the treatment of fault data also needs to be harmonized. Due to the absence of 3D fault surfaces in GTA3D it only might be possible to harmonize the main faults or fault systems, which have a sufficient horizontal length or a big vertical displacement (see Figure 12 and Figure 13).




Figure 12: few fault zones crossing the German-Dutch border (from DGM-NNL)




Figure 13: Depth maps of the base of the Upper North Sea Group and the base of the Lower Germanic Trias Group which clearly show that a few faults are crossing the German-Dutch border in the pilot area



## 4 LIST OF REFERENCES

- Baldschuhn, R., Binot, F., Fleig, S. & Kockel, F. (2001): Geotektonischer Atlas von Nordwest-Deutschland und dem deutschen Nordsee-Sektor. – Geol. Jb., A 153: S. 95; Stuttgart (Schweizerbart).
- Baldschuhn, R. F. (1996). Geotektonischer Atlas von Nordwestdeutschland. - 17 Teile, Kt., Taf. Hannover: BGR.
- Bombien et al. (2012) Der Geotektonische Atlas von Niedersachsen und dem deutschen Nordseesektor als geologisches 3D-Modell. In: Geowissenschaftliche Mitteilungen : GMit / Berufsverband Deutscher Geowissenschaftler.- 48 (2012), Seite 6-13.
- Doornenbal, J.C., Abbink, O.A., Duin, E.J.T., Dusar, M., Hoth, P., Jasionowski, M., Lott, G.K., Mathiesen, A., Papiernik, B., Peryt, T.M., Veldkamp, J.G. & Wirth, H., 2010. Introduction, stratigraphic framework and mapping. In: Doornenbal, J.C. and Stevenson, A.G. (Eds): Petroleum Geological Atlas of the Southern Permian Basin Area. EAGE Publications b.v. (Houten): 1-9.
- Doornenbal, J.C. and Stevenson, A.G. (editors), 2010. Petroleum Geological Atlas of the Southern Permian Basin Area. EAGE Publications b.v. (Houten). ISBN 978-90-73781-61-0
- Eurostat (2018). recalled 19. 10 2018 at  
<https://ec.europa.eu/eurostat/web/gisco/geodata/reference-data/administrative-units-statistical-units/countries#countries16>
- Helms, M (2018). The Geological 3D-Model for Lower Saxony. Presented at Workpackage Meeting 1 in Hannover; unpublished
- Ad-Hoc AG Hydrogeologie (2008). Sachstandsbericht für einen bundeseinheitlichen Produktkatalog zur wirtschaftlichen Anwendung oberflächennaher geothermischer Anlagen.  
recalled 05. 10 2018 at  
[http://www.infogeobasis.de/dokumente/download\\_pool/PKOG\\_Abschlussbericht\\_1.3\\_08-04-25.pdf](http://www.infogeobasis.de/dokumente/download_pool/PKOG_Abschlussbericht_1.3_08-04-25.pdf)
- Kombrink, H., Doornenbal, J. C., Duin, E. J., den Bulk, M., van Gessel, S., ten Veen, J. H., & Witmans, N. (2012). New insights into the geological structure of the Netherlands; results of a detailed mapping project. Netherlands Journal of Geosciences - Geologie en Mijnbouw, 91-4, S. 419-446.
- LBEG (2008). Erläuterung zum Geotektonischen Atlas als 3D-Modell für Niedersachsen. recalled 04. 10 2018 at  
<http://nibis.lbeg.de/cardomap3/project/cm3/Erlaeuterungstexte/Erl%C3%A4uterung%20zum%20GTA-3D.pdf>
- LBEG. (2018). <https://nibis.lbeg.de/cardomap3/>. recalled 05. 10 2018 at  
<https://nibis.lbeg.de/cardomap3/>
- Menning, M. (1997): Geologische Zeitskala der Mark Brandenburg. - 1 Bl., Geoforschungszentrum Potsdam NE-Deutschland, Brandenburg - - Altersbestimmungen -
- NLfB (1995). Karte der Lage der Quartärbasis in Niedersachsen und Bremen 1 : 500.000. –. Hannover.
- Ten Veen, J.H., Kruisselbrink, A.F., den Dulk, M. and Witmans, N., in prep. Digital Geological Model Noord Nederland (DGM-NNL), TNO report, xx pp.
- TNO (2018). DINOLoket. recalled 04. 10 2018 at <https://www.dinoloket.nl/table>
- VDI (2010). Richtlinienreihe VDI 4640 "Thermische Nutzung des Untergrunds". Düsseldorf.